

False vacuum decay via Coleman - de Luccia instanton in the theory with quasi-exponential potential

Michal Demetrian*

*Department of Theoretical Physics, Comenius University
Mlynská Dolina F2, 842 48, Bratislava, Slovak Republic*

(Dated: February 7, 2008)

The necessary and sufficient condition for the false vacuum decay in a de Sitter universe via Coleman - de Luccia instanton is applied on the scalar field with quasi-exponential potential.

PACS numbers: 98.80.Cq

I. INTRODUCTION

The theory of false vacuum decay in de Sitter universe via formation of rapidly expanding bubbles was introduced by Coleman and de Luccia [1]. The process has been studied intensively as an ingredient of old inflation [2] and of open inflation [3, 4]. This mode of vacuum decay is described by the Coleman - de Luccia (CdL) instanton - the nontrivial finite-action $O(4)$ solution to the Euclidean equations for coupled scalar and gravitational fields, with an effective potential containing a true vacuum (at which it has the global minimum equal to zero) and a false vacuum (at which it has a local minimum) separated from the true vacuum by a finite potential barrier. It is given by two functions $\Phi(\tau)$ and $a(\tau)$, where Φ is the scalar field, a is the radius of the three-spheres of homogeneity determined from the circumference and τ is the radius of the three-spheres of homogeneity. Denote the effective potential by V . Functions Φ and a obey equations

$$\ddot{\Phi} + \frac{3\dot{a}}{a} = V', \quad \dot{a}^2 = 1 + \frac{8\pi}{3} \left(\frac{1}{2}\dot{\Phi}^2 - V \right) a^2, \quad (1)$$

where the overdot denotes differentiation with respect to τ and the prime differentiation with respect to Φ . The boundary conditions ensuring finiteness of the (Euclidean) action

$$S_E = \int d^4x \sqrt{|g_E|} \left[\frac{1}{2} g_E(\partial\Phi, \partial\Phi) + V(\Phi) - \frac{R}{16\pi} \right]$$

are

$$a(0) = \dot{\Phi}(0) = \dot{\Phi}(\tau_f) = 0, \quad (2)$$

where $\tau_f > 0$ is defined by the equation $a(\tau_f) = 0$. The existence of a (nontrivial) solution of the system of equations (1,2) remained unclear for many years. It has been supposed that the bubble (if it exists) must fit into the sphere, which is the false vacuum (constant Φ) solution of the equations (1); and this statement has been translated approximatively into the condition that the curvature (absolute value of the second derivative) of the

potential V in the false vacuum must be bigger than $4 \times$ corresponding Hubble parameter squared.

II. THE NECESSARY AND SUFFICIENT CONDITION FOR THE EXISTENCE OF CDL INSTANTON

Being motivated by the arguments of the papers [5] and [7], the authors of the paper [8] have found a necessary condition and a sufficient condition for the existence of the CdL instanton. Before we formulate these conditions let us introduce useful notations used in what follows: by $V_M = V(\Phi_M)$ we denote the local maximum of V (the top of the barrier), and by $H(\Phi)$ the value of the Hubble parameter corresponding to the energy density equal to $V(\Phi)$, $H(\Phi) = \sqrt{8\pi V(\Phi)/3}$. The conditions read:

If the inequality $-V''_M/H_M^2 > 4$ holds then the CdL instanton exists.

If the CdL instanton exists then the inequality $-V''(\Phi)/H^2(\Phi) > 4$ holds somewhere in the barrier.

For a given theory (potential) the conditions leave a *gray zone* in the parameter space of the potential in which one cannot decide whether the CdL instanton exists or not.

III. QUASI-EXPONENTIAL POTENTIAL

We are interested in the theory with the potential

$$V(\Phi) = \left[V_0 + \frac{1}{2}\Phi^2 \right] e^{-\Phi/\Phi_0}, \quad (3)$$

where V_0 and Φ_0 are positive parameters. The classical evolution of the scalar field coupled to gravity and self-interacting by the potential similar to ours has been discussed in many works; as an example and source of further information we refer to the paper [9]. In order to apply our conditions to the theory (3) we have to do nothing but investigate elementary properties of the function $V(\Phi)$. The barrier exists only if $V_0 < 1/2\Phi_0^2$. If we

*Electronic address: demetrian@fmph.uniba.sk

introduce the new parameter

$$\Delta = \sqrt{1 - 2\frac{V_0}{\Phi_0^2}}, \quad \Delta \in [0, 1],$$

the formula (3) becomes

$$V(\Phi) = \frac{1}{2} [\Phi^2 + \Phi_0^2(1 - \Delta^2)] e^{-\Phi/\Phi_0}.$$

The local extremes of the function V are reached at the points

$$\Phi_{m,M} = \Phi_0(1 \mp \Delta) \text{ and } V_M = \Phi_0^2(1 + \Delta)e^{-1-\Delta}.$$

The sufficient condition holds only if

$$\frac{32\pi}{3}\Phi_0^2 < \frac{\Delta}{\Delta + 1}. \quad (4)$$

In words, the sufficient condition holds if the barrier is sufficiently thin. The parameter Δ measures the thickness of the barrier because, if Φ_+ and Φ_- are the inflexion points of V to the right and left of Φ_M , we have $\Phi_+ - \Phi_- = 2\Phi_0\sqrt{1 + \Delta^2}$. The necessary condition is fulfilled only if the curves $32\pi V(\Phi)/3$ and $-V''(\Phi)$ have at least one common point in the interval $[\Phi_-, \Phi_+]$ (The curves can have two intersection points in that interval). After a little bit more complicated algebra one gets that this is true only if

$$\frac{32\pi}{3}\Phi_0^2 < \frac{\sqrt{5 - 4\Delta^2} + \Delta^2 - 2}{1 - \Delta^2}. \quad (5)$$

The inequalities (4,5) are depicted in the figure 1.




FIG. 1: The blue (red) curve represents the sufficient (necessary) condition. The area between the curves is the gray zone for the potential (3).

Acknowledgments

The author thanks Vlado Balek for useful discussions. This paper was supported by the grant VEGA 1/0250/03.

-
- [1] S. Coleman and F. de Luccia, Phys. Rev. **D21**, 3305 (1980).
 - [2] A.H. Guth, Phys. Rev. **D23**, 347 (1981).
 - [3] M. Bucher, A.S. Goldhaber and N. Turok, Phys. Rev. **D52**, 3314 (1995).
 - [4] A.D. Linde and A. Mezhlumian, Phys. Rev. **D52**, 6789 (1995).
 - [5] S.W. Hawking and I.G. Moss, Phys. Lett. **110B**, 35 (1982).
 - [6] A.D. Linde, *Particle physics and inflationary cosmology*, Harwood, Chur, Switzerland (1990).
 - [7] L. Jensen and P.J. Steinhardt, Nucl. Phys. **B237**, 176 (1984).
 - [8] V. Balek and M. Demetrian, Phys.Rev. **D69**, 063518 (2004); gr-qc/0311040.
 - [9] P. Parsons and J.D. Barrow, Phys.Rev. **D51**, 6757 (1995); astro-ph/9501086.